

Figure 1 The DSC thermogram of polymorph A.

41.

5




Figure 2 The X-ray powder diffractogram of polymorph A.

5




Figure 3 The DSC thermogram of polymorph B

43

5




Figure 4 The X-ray powder diffractogram of polymorph B

5




Figure 5 The DSC thermogram of pseudopolymorph C1, a solvate of acetone

45

5




Figure 6 The TGA trace of pseudopolymorph C1, a solvate of acetone

46

5




Figure 7 The X-ray powder diffractogram of pseudopolymorph C1, a solvate of Acetone.

5




Figure 8 The DSC thermogram of pseudopolymorph C2, a solvate of 1-butanol

5




Figure 9 The TGA trace of pseudopolymorph C2, a solvate of 1-butanol

5




Figure 10 The X-ray powder diffractogram of pseudopolymorph C2, a solvate of 1-butanol

5




Figure 11 The DSC thermogram of pseudopolymorph C3, a solvate of 2-butanol

5




Figure 12 The TGA trace of pseudopolymorph C3, a solvate of 2-butanol

5




Figure 13 The X-ray powder diffractogram of pseudopolymorph C3, a solvate of 2-butanol

5




Figure 14 The DSC thermogram of pseudopolymorph C4, a solvate of 1,4-dioxane

5




Figure 15 The TGA trace of pseudopolymorph C4, a solvate of 1,4-dioxane.

55

5




Figure 16 The X-ray powder diffractogram of pseudopolymorph C4, a solvate of 1,4-dioxane

5




Figure 17 The DSC thermogram of pseudopolymorph C5, a solvate of methylacetate

5




Figure 18 The TGA trace of pseudopolymorph C5, a solvate of methylacetate

58

5




Figure 19 The X-ray powder diffractogram of pseudopolymorph C5, a solvate of methylacetate

5




Figure 20 The DSC thermogram of pseudopolymorph C6, a solvate of methylethylketone.

5




Figure 21 The TGA trace of pseudopolymorph C6, a solvate of methylethylketone

61

5




Figure 22 The X-ray powder diffractogram of pseudopolymorph C6, a solvate of methylethylketone

5




Figure 23 The DSC thermogram of pseudopolymorph C7, a solvate of THF

5




Figure 24 The TGA trace of pseudopolymorph C7, a solvate of THF

64

5




Figure 25 The X-ray powder diffractogram of pseudopolymorph C7, a solvate of THF

5




Figure 26 The DSC thermogram of pseudopolymorph C8, a solvate of toluene

5




Figure 27 The TGA trace of pseudopolymorph C8, a solvate of toluene

67

5




Figure 28 The X-ray powder diffractogram of pseudopolymorph C8, a solvate of toluene

5




Figure 29 The DSC thermogram of pseudopolymorph C9, a solvate of 1-pentanol

5




Figure 30 The TGA trace of pseudopolymorph C9, a solvate of 1-pentanol

70

5




Figure 31 The X-ray powder diffractogram of pseudopolymorph C9, a solvate of 1-pentanol

71

5




Figure 32 The DSC thermogram of polymorph D

72

5




Figure 33 The X-ray powder diffractogram of polymorph D

73

5




Figure 34 The powder X-ray diffractogram of a mixture of polymorphs A and B.

74

5




Figure 35 The powder X-ray diffractogram of a mixture of polymorph B and pseudo-polymorph C₁.

10